

Stressors Experienced during Deployment among Canadian Armed Forces Personnel:

Factor Structure of Two Combat Exposure Scales

Kerry Sudom, PhD*
Defence Research and Development Canada
Department of National Defence
101 Colonel By Drive
Ottawa, Ontario, Canada K1A 0K2
Kerry.Sudom@forces.gc.ca

Kimberley Watkins
Defence Research and Development Canada
Department of National Defence
101 Colonel By Drive
Ottawa, Ontario, Canada K1A 0K2
Kimberley.Watkins@forces.gc.ca

Jennifer Born
Defence Research and Development Canada
Department of National Defence
101 Colonel By Drive
Ottawa, Ontario, Canada K1A 0K2
Jennifer.Born@forces.gc.ca

Mark Zamorski, MD, MHSA
CF Health Services Group HQ
Department of National Defence
101 Colonel By Drive
Ottawa, Ontario, Canada K1A 0K2
Mark.Zamorski@forces.gc.ca

*Corresponding author

Abstract

Combat exposure is linked to increased mental health problems among military personnel. Reliable, precise, and efficient measurement of combat experiences can facilitate understanding of the effects of combat on mental health. This study examined the psychometric properties of two scales used to assess deployment experiences in Canadian Armed Forces (CAF) personnel in two different settings: during a third location decompression (TLD) program following deployment, and during routine post-deployment screening. Principal components analyses yielded similar results for the two measures. The components containing items relevant to Exposure to the Dead and Injured, Dangerous Environment, and Active Combat were common across both surveys. The TLD results revealed a fourth component, Personal Suffering, while the post-deployment screening data revealed a fourth component, Perceived Responsibility. This study found categories of deployment stressors that can be used for further analysis, and underscores the importance of assessing a broad range of exposures in the combat environment.

Stressors Experienced during Deployment among Canadian Armed Forces Personnel:
Factor Structure of Two Combat Exposure Scales

An abundance of research has documented the effects of exposure to military combat on psychological health problems, such as posttraumatic stress disorder (PTSD) and depression (e.g., Booth-Kewley, Larson, Highfill-McRoy, Garland, & Gaskin, 2010; Smith et al., 2008; Wells et al., 2010). Indeed, among military factors, combat exposure is typically the strongest and most consistent predictor of psychopathology (see Watkins, unpublished). Exposure to combat has also been found to be associated with behavioural outcomes such as aggression and problematic alcohol use (Bray, Brown, & Williams, 2013; Gallaway, Fink, Millikan, Mitchell, & Bell, 2013).

Understanding the effects of combat on these psychosocial outcomes demands reliable, precise, and efficient measurement of combat experiences. Such measurement is also necessary in order to control for different levels of combat exposure in observational studies focusing on other variables of interest, such as social support and leadership. A broad range of approaches has been used to this effect: some studies have used deployment to a particular theater of operation or to particular locations as a proxy for combat exposure (Boulos & Zamorski, 2013). Others have used a single, dichotomous self-report item for lifetime combat exposure (Sareen et al., 2007), and others have defined combat as endorsement of one or more of a limited list of specific experiences (e.g., Smith et al. 2008; Wells et al., 2010). Most recent research has focused on scales with a larger number (30 or more) of potential combat-related experiences, usually analyzed as a count of the total number of events experienced (e.g., Hoge et al., 2004).

This last approach is appealing in its simplicity and power, and clear dose-response relationships between the count of experiences and mental health during (Garber, Zamorski & Jetly, 2012) and after deployment (Hoge et al, 2004) have been documented. Fewer studies have systematically examined the effects of individual categories of exposure. It might be expected that certain combat experiences, such as handling bodies or seeing someone killed or injured, are likely to occur together rather than independently, in which case scales might be shortened to minimize respondent burden. Depending on the mission and on the role of particular groups of personnel, experiences may co-occur in different patterns or occur at different frequencies, suggesting that scales developed for one mission or one group may not apply to others. Finally, different types of combat exposures may have different effects on mental health outcomes.

Using factor analytic techniques, researchers have identified up to seven dimensions of combat exposure. Commonly found categories of deployment experiences include exposure to the combat environment (e.g., clearing homes or buildings), close physical engagement (e.g., engaging in hand-to-hand combat), and proximity to serious injury or death (e.g., handling or uncovering human remains) (Guyker et al., 2013). Other studies have found additional factors, such as sexual trauma, interpersonal distress, and personal injury (Katz, Cojucar, Davenport, Clarke, & Williams, 2012); exposure evoking emotion (e.g., knowing someone who was seriously injured/killed) (Gallaway et al., 2014); killing of the enemy (Killgore et al., 2008); and moral injury (violation of moral standards by self or others, Nash et al., 2013). As well, some investigators have captured dimensions reflecting stressful but typically non-traumatic deployment

experiences such as home front separation, exposure to a different culture, physical workload, boredom, and work frustration (Boermans, Kamhuis, Delahaij, Korteling, & Euwema, 2013; Waller et al., 2012).

Despite the profusion of studies, it is not clear which aspects are driving the observed similarities and differences in factor structure. Variations in items used, response categories, and type of deployment may partly explain the differences in findings. This paper uses data collected using two different combat exposure scales in two different populations of Canadian Armed Forces (CAF) personnel who deployed in support of the mission of Afghanistan. The objective is to explore similarities and differences in the factor structure of these two instruments. First, in Study 1, the factor structure of the most widely used combat exposure scale of dichotomous items (e.g., Guyker et al., 2013) was explored in a population of CAF personnel with diverse deployment experiences; this will provide insight into the extent to which factor structure depends on the diversity of deployment experiences in the deployed population, and whether the factor structure varies in the CAF compared to other militaries. Next, in Study 2, the factor structure of a frequency-based deployment experiences scale developed for peacekeeping operations is explored and compared with the structure identified in Study 1. Study 2 data were collected in an anonymous fashion during Third Location Decompression as opposed to in a confidential fashion during later post-deployment screening (Study 1), allowing for exploration into the extent to which these variables influence the factor structure. Ultimately, this work is expected to lead to more parsimonious and more complete measures of combat exposure and to exploration of the differential health effects of various dimensions of combat exposure.

Study 1: Combat Experiences Assessed during Enhanced Post-Deployment Screening

Method

Procedure

The Enhanced Post-Deployment Screening (EPDS) process is mandatory for all CAF members who have deployed overseas for 60 days or longer; it is administered between 90 and 180 days after returning to Canada. The EPDS involves a confidential, although not anonymous, paper-and-pencil health questionnaire followed by an interview with a mental health clinician. Components of the health questionnaire were analyzed in this study. The EPDS received ethics approval from Veritas IRB, Dorval, Quebec.

Participants

Participants were 14,616 CAF personnel who deployed in support of the later phases of the combat and peace-support mission in Afghanistan (Operation ATHENA) and completed the EPDS between 2009 and 2012. Personnel were largely deployed in Kandahar Province, however the average threat level in this sample was relatively low because it included some personnel deployed to safe air bases in the region and because the threat level in Kandahar declined significantly in 2010 and onward. Personnel in Kandahar fulfilled a broader range of roles than a US Army Brigade Combat Team, with similar proportions serving in remote locations (e.g., a checkpoint), semi-remote locations (e.g., a forward operating base), and “inside the wire” at Kandahar Airfield (Garber, Zamorski & Jetly, 2012). After removing 1013 cases (7%) that were missing exposure data using listwise deletion, the dataset contained 13,603 individuals.

The EPDS sample consisted of 90.8% males ($n=12,346$) and 9.2% females ($n=1,253$). At the time of post-deployment screening, over half (54.1%) of participants

were under the age of 32, and had been in the CAF for 10 or fewer years (59.6%). The majority were Regular Force (85.1% Regular Force, 14.9% Reservists), junior NCMs (8.4% Privates or equivalent; 58.8% other junior NCMs), and in the Army (82.7%), followed by Air Force (14.5%) and Navy (2.8%).

Measures

Combat exposure was assessed using a modified version of a widely-used 34-item Combat Experiences Scale (CES) developed by the Walter Reed Army Institute for Research. This scale was developed and validated in US Army Brigade Combat Teams, and assesses events experienced during deployment (e.g., “being attacked or ambushed,” “participating in improvised explosive device [IED]/mine clearing”). For the EPDS, four items of the original scale were removed because of concerns that a positive response might require investigation into potential misconduct (e.g., “witnessing mistreatment of a non-combatant”). The response options for each item were “yes” and “no,” for ever having been exposed during the most recent deployment.

Statistical Analyses

The factor structure was explored using principal components analysis (PCA), using an oblique (Promax) rotation to account for the likely correlation of the underlying factors, consistent with past research approaches (e.g., Guyker et al., 2013). Cross loading items were resolved using a combination of empirical (i.e., higher loading score) and subjective (i.e., thematic and theoretical relevance) criteria. Stata version 13.1 was used for analysis in order to calculate the polychoric correlations needed to perform the PCA for dichotomous items (Uebersax, 2006).

Results

As shown in Table 1, the most commonly reported combat exposure items were “Receiving incoming artillery, rocket or mortar fire” (64.3%), “Improvised explosive device (IED)/booby trap exploded near you” (61.3%), and “Having hostile reactions from civilians” (58.9%). The least common items, including “Engaging in hand-to-hand combat”, “Feeling responsible for the death a non-combatant” and “Feeling responsible for the death of Canadian or ally personnel,” were endorsed by fewer than three percent of respondents.

PCA was run on all 30 items. Several aspects of principal components analysis, including the eigenvalues exceeding 1.00 (Kaiser’s criterion) and visual interpretation of the scree plot and the pattern matrix, suggested that four factors should be retained. In combination, these four factors explained 81.9% of the variance. The first factor, labelled “Dangerous Environment,” reflected items that were related to risks in the deployed environment, including exposure to small arms fire, IEDs, and hostile civilians, and accounted for 63.8% of the variance. The second factor, labelled “Exposure to the Dead and Injured” reflected exposures to the aftermath of conflict involving death or serious injury of others; it accounted for 7.6% of the variance. The third factor, labelled “Active Combat,” represented exposure to armed combat and included issues such as shooting or directing fire at the enemy. This factor accounted for 5.7% of the variance. The final category, accounting for 4.8% of the variance, included three items: “Engaging in hand-to-hand combat”, “Feeling directly responsible for the death of a non-combatant”, and “Feeling directly responsible for the death of a Canadian or ally personnel.” These items were grouped under the category titled “Perceived Responsibility” because of the

potential moral implications of the events included in this factor, as they likely reflected unintentional deaths.

Eleven items cross-loaded in the rotated solution; these items were assigned to factors based on empirical and subjective criteria. For example, “Seeing a unit member blown up or burned alive” was assigned to the Exposure to the Dead and Injured factor because the item loaded much higher onto this factor. Alternatively, “Improvised explosive device (IED)/booby trap exploded near you” was assigned to Dangerous Environment, which contained similar IED-related items. Three items with a loading score of less than .4 or that did not improve the reliability of the component did not load on any of the factors: “Had a close call, was shot, hit but protective gear saved you” (reported by 10.3%); “Witnessing a friendly fire incident” (7.5%); and “Being wounded/injured” (7.6%). Table 1 shows the rotated pattern matrix for this analysis.

The four factors exhibited intercorrelations between .29 and .62 (Table 2). The highest correlation was between Dangerous Environment and Exposure to the Dead and Injured, while the lowest was between Dangerous Environment and Perceived Responsibility.

The first three factors, Dangerous Environment, Exposure to the Dead and Injured, and Active Combat, exhibited good internal consistency (KR-20 coefficients of .88, .78, and .70, respectively). However, the fourth factor, Perceived Responsibility, exhibited low reliability (.29).

Study 2: Combat Experiences Assessed during Third Location Decompression

Method

Procedure

The Third Location Decompression (TLD) program aims to assist military members in transitioning back to the civilian environment following a deployment through rest, recreation, and educational sessions in a neutral setting (Cyprus) before returning to Canada (Garber & Zamorski, 2012). Questions on combat exposure were included in an anonymous, voluntary, paper-and-pencil TLD evaluation questionnaire offered at the end of TLD.

Participants

Using listwise deletion of cases missing combat exposure data, the dataset contained 9,028 participants returning from a combat and peace support mission in Kandahar Province, Afghanistan. The TLD sample was 91.7% male ($n=8,157$) and 8.3% female ($n=737$). Over half (54.1%) were under the age of 32, and had 10 or fewer years of service (56.9%). The majority were Regular Force (84.9% Regular Force, 15.1% Reservists) and junior NCMs (18.2% Privates or equivalent; 50.6% other junior NCMs). The questionnaire did not include an item on element (Army, Navy, Air Force), but administrative data indicates that at least 90% were Army personnel.

Measures

Twenty items from the Stress in Military Service Questionnaire (SMSQ) from the CAF Human Dimensions of Operations study (Dobрева-Martinova, 1998) were used to assess deployment experiences. The SMSQ was developed to explore the stressors typical of the peacekeeping deployments in the 1990s, but includes combat-related experiences. Participants answered each item in terms of the frequency of encounters with various deployment experiences, such as “Being in armed combat,” “Seeing a CAF member die,”

and “Being sexually assaulted.” Response options were “Never,” “Once,” “A few times,” and “Regularly,” with higher scores indicating greater exposure to that experience.

Statistical Analysis

SPSS version 22 was used to explore the factor structure of the items, again using PCA with an oblique (Promax) rotation. Cross loading items were again resolved using a combination of empirical (higher loading score) and subjective criteria (thematic and theoretical relevance). Given that the study data came from anonymous, voluntary evaluation of an educational intervention, no Research Ethics Board approval was sought.

Results

As shown in Table 3, the most commonly reported combat exposure items were “Being subject to shelling and/or artillery/mortar fire” (82.1%), “Seeing serious injuries” (64.8%), and “Being fired at” (64.8%). Other items, including “Being held hostage” (3.0%) and “Being sexually assaulted” (2.6%), were endorsed by a much smaller percentage of participants.

The PCA, run using the general approach used in Study 1, supported retaining four factors. Items with a loading score of less than .4 and those that did not improve the reliability of the component were excluded from the factor structure, which resulted in the removal of one item (“Seeing physical violence”, reported by 56.9% of respondents).

The four factors exhibited intercorrelations between .30 and .60 (Table 4), justifying the use of an oblique rotation. The highest correlation was between Dangerous Environment and Active Combat; the lowest was between Personal Suffering and Active Combat.

The first factor, labelled “Exposure to the Dead and Injured,” reflected exposures to the consequences of combat events involving the death or serious injury of others, including seeing multiple deaths and handling bodies or body parts. This factor explained 37.4% of the variance. The second factor, labelled “Personal Suffering,” contained items reflecting personal harassment and threats such as physical assault or being threatened with assault. This factor explained 13.2% of the variance. The experiences that loaded onto this factor were much less commonly reported (3% to 15% of participants) than the items in the other factors (34% to 65% of participants). The third factor, labelled “Dangerous Environment,” contained items reflecting dangers in the combat environment that do not necessarily involve personal injury, such as dangerous training conditions. This factor explained 8.4% of the variance. Finally, the fourth factor, labelled “Active Combat,” reflected exposure to traditional combat events such as being fired at or harming another person. This factor explained 5.2% of the variance, for a total explained variance of all four factors of 64.1%. Inter-item reliabilities for each component were in the acceptable range, with Cronbach’s alphas of .88, .78, .82, and .78, for Exposure to the Dead and Injured, Personal Suffering, Dangerous Environment, and Active Combat, respectively. The rotated pattern matrix for the items is shown in Table 3.

Discussion

Summary of Key Findings

Exploratory factor analyses of two different combat exposure scales each yielded four factors, three of which appeared very similar in the two studies (“Dangerous Environment,” “Exposure to the Dead and Injured,” and “Active Combat.”). The

commonality of the factor structure was seen despite important differences between 1) the context for which the scale was developed (combat vs. peacekeeping operations); 2) somewhat different items; 3) the response categories (use of dichotomous vs. frequency); 4) the context and timing of the assessment (during post-deployment screening three to six months after return vs. during TLD on the way home from deployment; 5) the anonymity of the assessment; 6) the diversity of the study population (a more diverse group, including personnel deployed to a safe air base vs. personnel deployed to a high-threat area); and 7) the average threat level of the population (a lower threat period late in the conflict vs. a higher threat period early in the conflict).

One unique factor emerged in each of the studies: In Study 1, a fourth factor, labelled “Perceived Responsibility” came out, while in Study 2, a different fourth factor, labelled “Personal Suffering” was identified. The internal consistency values of all of the components were acceptable, with the exception of Perceived Responsibility (Study 1), which contained only three items. The analyses revealed several items (three in Study 1 and one in Study 2) which did not load on any of the factors.

Comparison with Previous Research

These results show both commonalities and discrepancies with three US factor analytic studies that used different versions of the same measure used in Study 1. Kilgore et al. (2008), Guyker et al. (2013), and Gallaway et al. (2014) identified seven, four, and three factors, respectively. For example, Kilgore et al.’s first factor, labelled “violent combat exposure,” included items that in Study 1 loaded on “dangerous environment” and others that loaded on “exposure to the dead and injured.” Gallaway et al.’s (2014) “active exposure” factor had strongly-loading items on all four of Study 1’s factors.

Similar discrepancies are seen between the factor structure of Study 1 and those of Guyker et al. (2013).

Some of these differences are likely due to technical differences in the studies, notably the use of dichotomous vs. frequency-based response categories, the extraction method (principal axis factoring vs. PCA), the particular items included in the studies, and the use or non-use of polychoric correlations in the extraction process for dichotomous items. Other differences may relate to decisions as to which factor to assign weakly-loading or cross-loading items. The principal differences in the observed factor structure seem readily attributable to differences in the items. For items common to both Study 1 and Study 2, the factor structures appear to be conceptually coherent.

A possible explanation for the coherent factor structure in Study 1 and Study 2 and the divergent factor structure between Study 1 and that seen in US studies with the same items (or subsets thereof) is that the underlying experiences occur together in ways that are idiosyncratic to the study population, its role on deployment, and the circumstances on the ground at the time. Consistent with this hypothesis, as opposed to technical issues or limitations associated with factor analysis, Gallaway et al. (2014) found convergent findings in both exploratory factor analysis and confirmatory factor analysis in their study population.

Limitations

Exploratory factor analysis has intrinsic limitations, related in part to the broad range of choices investigators have with respect to the extraction method, use of correlation vs. covariance matrices, rotation options, and the many subjective choices surrounding issues such as cross-loading items and cut-offs for eigenvalues.

Confirmatory factor analysis is a potential solution, but it is limited in applications without a well thought-out conceptual framework or well-validated factor structure to draw upon. We chose to use an exploratory approach in both studies precisely because of the differences in factor structure observed in other studies.

The data collected in both studies are based on self-reported, retrospective recall of events during deployment. Self-reported combat exposure has the potential for biased or inaccurate recall, particularly in the case of post-deployment screening, which occurs several months after returning from a deployment (Richardson, Frueh, & Acierno, 2010). In addition, although confidentiality is assured, some respondents may be reluctant to admit involvement in certain events, particularly those involving perceived responsibility for the death of others, due to fear of personal or career repercussions.

The items falling into the Perceived Responsibility component in Study 1 yielded a subscale containing only three items, which exhibited low reliability. Having few items on a subscale is commonly associated with low reliability (Cortina, 1993). Although one of these items, involvement in hand-to-hand combat, may seem conceptually unrelated to the other items on perceived responsibility for unintended death, this item loaded with others in at least one other study (Killgore et al., 2008), suggesting that this may not be a spurious or purely technical finding. As well, qualitative research on killing in hand-to-hand combat among military personnel shows that this experience may have particularly strong psychological repercussions (Jensen & Simpson, 2014), suggesting that it may be associated with perceived responsibility for unintended death in at least some who have had this experience. Future research is needed to further develop this subscale to more accurately assess the experiences of military personnel. Qualitative research to determine

the events involving personal responsibility may yield further insight into this category of stressors, and allow for the development of a scale to fully capture such experiences during deployment. Although such scales have been developed for other militaries (e.g., Nash et al., 2013), it is important that scales capture the experiences of CAF members as well as the current types of conflict, and continue to be updated as involvement in missions changes.

Combat exposure events were assessed in dichotomous response format during post-deployment screening. The creation of exposure categories does not account for the relative degree of stress associated with each experience. Certain types of exposures may be more likely to lead to the development of psychological health problems. As well, there may be subjectivity in reporting of certain events. This is especially important for some items, such as feeling responsible for the death of another individual, in which there is particular room for subjectivity or errors in recall. In contrast, items such as being under mortar attack or IED blast are more objective, and the dichotomous response categories are more likely to accurately reflect participants' experiences. However, despite its limitations, use of a dichotomous response format for combat exposure items may reduce the reporting bias that can occur for questions about frequency, duration, or intensity of stress (Lund, Foy, Sippelle, & Stratchan, 1984).

Implications

In the context of the present paper, the factor structure of deployment experiences provides insight into which experiences tend to occur together in a given individual in the study populations. Study 1 and Study 2 showed convergent results, suggesting that, in the same region over different time periods, the experiences of military personnel

clustered together in reproducible ways. Where differences were seen, they could be traced to the disparities in items on the two surveys.

A parsimonious explanation for the divergent results between the factor structure identified in the current study and the structures found in other populations at other times is that the pattern of shared experiences happened to be different for that group at that time. The larger size and greater diversity of personnel in both of our studies relative to similar work may have facilitated convergence in the factor structure, in that the role of particular shared experiences in, for example, a single Brigade Combat Team, may have been lessened.

That certain experiences happened to occur together should not be interpreted as evidence of homogeneous impacts on mental health. The motivation for this study was largely technical in nature: to inform the development of a more compact exposure scale for the purpose of controlling for combat exposure in epidemiological studies and for exploring the effects of different types of combat exposure. Before item reduction, the association of each of those experiences with mental health problems will need to be explored. Other recent work on the psychosocial consequences of different types of combat exposure (Gallaway et al., 2014; Killgore et al., 2008) has not explicitly done so.

The presence of unique factors (Perceived Responsibility in Study 1 and Personal Suffering in Study 2) points to the need for greater conceptual and empirical clarity. Developing more items (or better items) in these two areas will help to understand if they contribute to the burden of post-deployment mental disorders. Qualitative research in these domains will be helpful. In addition, qualitative research on cross-loading items

may provide a deeper understanding of the nature of each of the identified factors and why certain experiences tended to occur together.

Four items of the original CES in Study 1 were removed due to concerns that they may require investigation into potential misconduct. Inclusion of these items (e.g., witnessing mistreatment of non-combatants) may have yielded a more statistically reliable perceived responsibility scale, or a separate factor pertaining to violations of ethical norms. As well, ethical violations have conceptual and empirical links to mental health, underlining the importance of work in this area.

Two other types of deployment experiences should also be explored in conjunction with combat experiences in future work to provide the fullest possible picture of how those experiences affect deployed personnel. Non-combat stressors (such as exposure to a harsh environment, separation from family, lack of physical intimacy) may exert independent negative effects or may moderate the effects of traumatic experiences; these effects may differ in-theater compared to post-deployment. Positive deployment experiences (e.g., gratitude from civilians, camaraderie) raise similar questions. Positive mental health may both serve as a stress buffer and contribute independently to well-being; it has been targeted in resilience training programs, particularly in the US Army. Understanding how all of these factors (traumatic stressors, non-traumatic stressors, and positive experiences) influence health may identify targets for intervention. Non-combat stressors and positive experiences are appealing targets in that the military may have greater control over them than traumatic combat stressors. Finally, exploration of all of these experiences in future non-combat operations is essential.

Conclusion

It is important to systematically assess combat exposure in order to develop valid and reliable measures that accurately reflect the experiences of military personnel. Accurate information about traumatic experiences, both traditional combat events, such as being involved in a firefight, as well as events involving perceived responsibility for harm of others, can provide information to clinicians to help focus assessment and treatment strategies for mental health issues. The current study found categories of deployment stressors that can be used for further analysis.

The similarities and differences in the factor structure of similar combat exposure scales in different groups raises the possibility that combat exposures cluster somewhat idiosyncratically, particularly in smaller groups that work together in a particular place at a particular time. Additional exposure items are needed to capture dimensions of negative deployment experiences (e.g., moral injury, Nash et al., 2013) that are not well-covered by the most widely-used scales. Also, investigators should consider the possibility that experiences that happen to occur together could have very different impacts on mental health. Finally, exploration of the health effects of traumatic deployment experiences needs to occur in conjunction with exploration of the effects of non-combat stressors and positive deployment experiences.

References

Boermans, S. M., Kamhuis, W., Delahaij, R., Korteling, J. E., & Euwema, M. C. (2013). Perceived demands during military operations. *Military Medicine, 178*, 722-728.

Booth-Kewley, S., Larson, G. E., Highfill-McRoy R. M., Garland, C. F., & Gaskin, T. A. (2010). Correlates of posttraumatic stress disorder symptoms in Marines back from war. *Journal of Traumatic Stress, 23*, 69-77.

Boulos, D. L., & Zamorski, M. A. (2013). Deployment-related mental disorders in Canadian Forces personnel deployed in support of the mission in Afghanistan, 2001 - 2008. *Canadian Medical Association Journal, 185*, E545-E552. Retrieved from doi: 10.1503/cmaj.122120

Bray, R. M., Brown, J. M., Williams, J. (2013). Trends in binge and heavy drinking, alcohol-related problems, and combat exposure in the U.S. military. *Substance Use and Misuse, 48*, 799-810.

Cortina, J. M. (1993). What is coefficient alpha? An examination of theory and applications. *Journal of Applied Psychology, 78*, 98-104.

Dobрева-Martinova, T. (1998). *Psychometric properties of the Stress in Military Service Questionnaire based on surveys of deployed Canadian Forces personnel*. DHRRE Sponsor Research Report 98-16. Ottawa, Canada: Director Human Resources Research and Evaluation.

Gallaway, M. S., Fink, D. S., Millikan, A. M., Mitchell, M. M., & Bell, M. R. (2013). The association between combat exposure and negative behavioral and psychiatric conditions. *Journal of Nervous and Mental Disease, 201*, 572-578.

Gallaway, M. S., Mitchell, M., Fink, D. S., Forsys-Donahue, K., Pecko, J., Bell, M. R., & Millikan, A. M. (2014). Combat exposure factors associated with the likelihood of behavioral and psychiatric issues. *Military Behavioral Health*, available online at <http://www.tandfonline.com/doi/full/10.1080/21635781.2014.897205>.

Garber, B. G., & Zamorski, M. A. (2012). Evaluation of a third-location decompression program for Canadian Forces members returning from Afghanistan. *Military Medicine, 177*, 397-403.

Garber, B. G., Zamorski, M. A., & Jetly, R. (2012). Mental health of Canadian Forces personnel while on deployment to Afghanistan. *Canadian Journal of Psychiatry, 57*, 736-744.

Guyker, W. M., Donnelly, K., Donnelly, J. P., Dunnam, M., Warner, G. C., Kittleson, J., Bradshaw, C. B., Alt, M., & Meier, S. T. (2013). Dimensionality, reliability, and validity of the Combat Experiences Scale. *Military Medicine, 178*, 377-384.

Hoge, C. W., Castro, C. A., Messer, S. C., McGurk, D., Cotting, D. I., & Koffman, R. L. (2004). Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *New England Journal of Medicine, 351*, 13-22.

Jensen, P. R., & Simpson, D. (2014). A qualitative analysis of the experience and impact of killing in hand-to-hand combat. *Journal of Traumatic Stress, 27*, 468-473.

Katz, L. S., Cojucar, G., Davenport, C., Clarke, S., & Williams, J. C. (2012). War Experiences Inventory: Initial psychometric and structural properties. *Military Psychology, 24*, 48-70.

Killgore, W. D., Cotting, D. I., Thomas, J. L., Cox, A. L., McGurk, D., Vo, A. H., Castro, C. A., & Hoge, C. W. (2008). Post-combat invincibility: violent combat experiences are associated with increased risk-taking propensity following deployment. *Journal of Psychiatric Research, 42*, 1112-1121.

Lund, M., Foy, D., Sippelle, C., & Strachan, A. (1984). The Combat Exposure Scale: A systematic assessment of trauma in the Vietnam War. *Journal of Clinical Psychology, 40*, 1323-1328.

Nash, W. P., Marino Carper, T. L., Mills, M. A., Au, T., Goldsmith, A., & Litz, B. T. (2013). Psychometric evaluation of the Moral Injury Events Scale. *Military Medicine*, *178*, 646-652.

Richardson, L. K., Frueh, B. C., & Acierno, R. (2010). Prevalence estimates of combat-related PTSD: a critical review. *Australian and New Zealand Journal of Psychiatry*, *44*, 4-19.

Sareen, J., Cox, B. J., Afifi, T. O., Stein, M. B., Belik, S. L., Meadows, G. et al. (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care: findings from a large representative sample of military personnel. *Archives of General Psychiatry*, *64*, 843-852.

Smith, T. C., Ryan, M. A. K., Wingard, D. L., Slymen, D. J., Sallis, J. F., & Kritzer-Silverstein, D. (2008). New onset and persistent symptoms of post-traumatic stress disorder self reported after deployment and combat exposures: prospective population based US military cohort study. *British Medical Journal*, *336*, 366-371.

Uebersax, J. S. (2006). The tetrachoric and polychoric correlation coefficients. Statistical Methods for Rater Agreement web site. Available at: <http://john-uebersax.com/stat/tetra.htm>. Accessed October 16, 2003.

Waller, M., Treloar, S. A., Sim, M. R., McFarlane, A. C., McGuire, A. C. L., Bleier, J., & Dobson, A. J. (2012). Traumatic events, other operational stressors and physical and mental health reported by Australian Defence Force personnel following peacekeeping and war-like deployments. *BMC Psychiatry*, *12*: 88. Available online at <http://www.biomedcentral.com/1471-244X/12/88>.

Watkins, K. (2014). *Deployment stressors: A review of the literature and implications for members of the Canadian Armed Forces*. Manuscript submitted for publication.

Wells, T. S., LeardMann, C. A., Fortuna, S. O., Smith, B., Smith, T. C., Ryan, M. A. K., Boyko, E. J., Blazer, D. (2010). A prospective study of depression following combat deployment in support of the wars in Iraq and Afghanistan. *American Journal of Public Health, 100*, 90-99.

Table 1: Factor Loadings based on a Principal Components Analysis using for Deployment Experiences measured during Post-Deployment Screening¹

Exposure category	Item	Proportion who experienced at least once	Factor loadings			
			1	2	3	4
Dangerous Environment (KR-20 = .88)	Having hostile reactions from civilians	58.9%	1.02			
	Clearing/searching homes or building	29.6%	0.91			
	Participating in IED/mine clearing	29.2%	0.91			
	Clearing/searching caves or bunkers	11.7%	0.80			
	Working in areas that were mined or had IEDs	39.8%	0.80			
	Witnessing an accident which resulted in serious injury or death	38.0%	0.77			
	Being in threatening situation where you were unable to respond because of rules of engagement	20.1%	0.64			
	Receiving small arms fire	42.7%	0.60		0.44	
	Having difficulty distinguishing between combatants and non-combatants	21.5%	0.60			
	Being attacked or ambushed	43.0%	0.51		0.43	
	Improvised explosive device (IED)/booby trap exploded near you	61.3%	0.44	0.44		
	Seeing ill/injured women or children who you were unable to help	24.5%	0.42			
	Exposure to the Dead and Injured (KR-20 = .78)	Knowing someone seriously injured or killed	35.3%		0.85	
Handling or uncovering human remains		20.0%		0.80		
Seeing dead bodies or human remains		46.1%		0.69		
Seeing a unit member blown up or burned alive		12.0%	0.34	0.63		
Having a member of your own unit become a casualty		39.8%	0.36	0.45		
Had a buddy shot or hit who was near you		10.4%	0.37	0.40		

¹ Factor loadings of less than .3 are suppressed in this table.

	Seeing dead or injured Canadians	34.7%	0.37	0.37		
Active Combat (KR-20 = .70)	Feeling directly responsible for the death of an enemy combatant	10.4%			0.85	
	Shooting or directing fire at the enemy	28.7%	0.47		0.63	
	Calling in fire on the enemy	13.8%			0.62	
	Receiving incoming artillery, rocket or mortar fire	64.3%			0.58	
	Sniper fire	6.9%	0.39		0.47	
Perceived Responsibility (KR-20 = .29)	Engaging in hand-to-hand combat	0.9%				0.78
	Feeling directly responsible for the death of a non-combatant	1.9%			0.30	0.73
	Feeling directly responsible for the death of a Canadian or ally personnel	2.6%		0.30		0.53
	% of variance		63.8%	7.6%	5.7%	4.8%

Table 2: Component Correlations: Post-Deployment Screening Data

Factor	Dangerous Environment	Aftermath of Conflict	Active Combat	Perceived Responsibility
Dangerous Environment	1			
Exposure to the Dead and Injured	0.62	1		
Active Combat	0.55	0.47	1	
Perceived Responsibility	0.29	0.31	0.36	1

Table 3: Factor Loadings based on a Principal Components Analysis for Deployment Experiences measured during Third Location Decompression¹

Exposure Category		Proportion who experienced at least once	Factor Loadings			
			1	2	3	4
Exposure to the Dead and Injured ($\alpha = .88$)	Seeing multiple deaths	43.4%	.90			
	Handling bodies or body parts	38.1%	.90			
	Seeing a non-military person die	46.5%	.85			
	Seeing serious injuries	64.8%	.78			
	Seeing a CAF member die	34.4%	.77			
	Seeing physical violence	56.1%	.37			.31
Personal Suffering ($\alpha = .78$)	Being sexually assaulted	2.6%		.91		
	Being held hostage/captive	3.0%		.89		
	Being physically assaulted	5.1%		.84		
	Being seriously injured	7.6%		.74		
	Being threatened with assault	15.3%		.45	.41	
Dangerous Environment ($\alpha = .82$)	Dangerous traffic conditions/incidents	63.6%			.90	
	Dangerous training conditions/incidents	44.0%			.88	
	Seeing widespread suffering	52.2%	.41		.63	
	Seeing widespread destruction	61.4%			.58	
	Being threatened with death	42.8%			.49	
Active Combat ($\alpha = .78$)	Being in armed combat	59.6%				.85
	Being fired at	64.8%				.84
	Being subject to shelling and/or artillery/mortar fire	82.1%				.72
	Harming a person	39.3%				.67
	% of variance		37.4%	13.2%	8.4%	5.2%

¹ Factor loadings of less than .3 are suppressed in this table.

Table 4: Component Correlations: Third Location Decompression Data

Factor	Aftermath of Conflict	Personal Suffering	Dangerous Environment	Active Combat
Exposure to the Dead and Injured	1			
Personal Suffering	.33	1		
Dangerous Environment	.52	.35	1	
Active Combat	.51	.30	.60	1

Note: All correlations are significant at the $p < .01$ level.